## **MSS Prenatal Eligibility Tool**

07/01/09

## **Instructions:**

- An \* asterisk indicates MSS targeted risk factors that need a clinician to determine the clients risk criteria (A, B or C).
- After screening the client for the MSS targeted risk factors, document the date(s) in the appropriate A, B or C column for any identified criteria, sign the last page noting who made the determination and assign the level of service.

| m                          |   |   |   |  |  |
|----------------------------|---|---|---|--|--|
| Targeted<br>Risk<br>Factor | A | В | C | Risk Factor Criteria   |  |
| Race                       |   |   |   | <b>B.</b> American Indian, Alaska Native or non-Spanish speaking indigenous women from the Americas (e.g. women whose primary language is Mixteco, Mam, or Kanjobal, etc.) |  |
|                            |   |   |   | C. African American or Black   |  |
| Prenatal<br>Care           |   |   |   | A. Greater than or equal to (≥) 14 and less than (<) 24 weeks gestation and no prenatal care started at the time of screening.   |  |
|                            |   |   |   | B. Greater than or equal to (≥) 24 weeks gestation before prenatal care was started  |  |
|                            |   |   |   | <b>B.</b> Greater than or equal to (≥) <b>24</b> weeks gestation <b>and no prenatal care</b> has started.  |  |
| Nutrition                  |   |   |   | Food Insecurity:  A. Runs out of food before the end of the month or cuts down on the amount eaten to feed others  |  |
|                            |   |   |   | <ul> <li>Pre-pregnancy BMI:</li> <li>*A. Pre-pregnancy BMI less than (&lt;) 18.5 and adequate weight gain based on Institute of Medicine guidelines.</li> </ul>            |  |
|                            |   |   |   | *C. Pre-pregnancy BMI less than (<) 18.5 and inadequate weight gain based on Institute of Medicine guidelines.   |  |
|                            |   |   |   | A. Pre-pregnancy BMI 25.0 to 29.9  |  |
|                            |   |   |   | *A. Pre-pregnancy BMI greater than or equal to (≥) 30 and adequate weight gain based on Institute of Medicine guidelines.  |  |
|                            |   |   |   | *B. Pre-pregnancy BMI greater than or equal to (≥) 30 and weight gain outside of the Institute of Medicine guidelines.   |  |
| Medical                    |   |   |   | Inter-pregnancy interval: A. Current pregnancy conception less than (<) 9 months from the end of the last pregnancy  |  |
|                            |   |   |   | Diabetes:  B. History gestational diabetes with last pregnancy   |  |
|                            |   |   |   | C. Pre-existing Diabetes- type 1 or 2  |  |
|                            |   |   |   | C. Current gestational diabetes  |  |
|                            |   |   |   | Multiples: C. Currently pregnant with multiples (2 or more babies)   |  |
|                            |   |   |   | Hypertension/Gestational Hypertension:   |  |
|                            |   |   |   | A. Gestational Hypertension in past pregnancy  |  |
|                            |   |   |   | C. Chronic Hypertension: Hypertension diagnosed prior to pregnancy or before 20 weeks gestation  |  |
|                            |   |   |   | C. Current pregnancy induced hypertension (gestational hypertension) starting greater than 20 weeks gestation  |  |
|                            |   |   |   | Low Birth Weight (LBW) or Preterm birth/labor/fetal death:   |  |
|                            |   |   |   | <b>C.</b> Prior LBW infant (less than 5lb 8 oz) <b>and/or</b> premature infant (less than 37 weeks); Prior fetal death (fetus greater than 20 weeks gestation).            |  |
|                            |   |   |   | C. Current pregnancy - diagnosed with <b>preterm labor</b> during this pregnancy or is on treatment or bed rest to prevent preterm birth.                                  |  |

| Risk<br>Factor                   | A  | В                  | C   | Risk Factor Criteria   |  |  |
|----------------------------------|--|--------------------|---|--|--|--|
| Maternal                         |  |                    |   | A. 17 years of age at conception   |  |  |
| Age                              |  |                    |   | <b>B.</b> Less than or equal to (≤) 16 years of age at conception  |  |  |
|                                  |  |                    |   | <b>A.</b> Greater than or equal to (≥) <b>35</b> years of age at conception <b>and</b> this is not her first pregnancy <b>and</b> she did not use assisted reproductive technology (ART) for this pregnancy  |  |  |
|                                  |  |                    |   | <ul> <li>B. Greater than or equal to (≥) 35 years of age at conception and one of the following:</li> <li>(1) First pregnancy</li> <li>(2) Current pregnancy via assisted reproductive technology (ART)</li> </ul>   |  |  |
| Maternal                         |  |                    |   | A. Quit smoking and/or or using tobacco prior to pregnancy or upon diagnosis of pregnancy  |  |  |
| Tobacco                          | ohacco   |                    |   |  |  |  |
| Use                              |  |                    |   | <b>B.</b> Smokes and/or uses tobacco during pregnancy  |  |  |
| Alcohol &<br>Substance           |  |                    |   | *B. Actively engaged in alcohol/drug treatment program and has not used for greater than or equal to (≥) 90 days.  |  |  |
| Abuse or                         |  |                    |   | *B. Stopped use of alcohol (see clarification table), illicit substances, or non-prescriptive  |  |  |
| Addiction                        |  |                    |   | use of prescriptive drugs <b>following pregnancy diagnosis and</b> has not used for more than or equal to $(\geq)$ <b>90 days</b>  |  |  |
|                                  |  |                    |   | *C. Stopped use of alcohol ( see clarification notes), illicit substances, or non-prescriptive   |  |  |
|                                  |  |                    |   | use of prescriptive drugs <b>following pregnancy diagnosis an d</b> has not used for less than (<90)   |  |  |
|                                  |  |                    |   | *C. Any use of alcohol, illicit substances, or non-prescriptive use of prescriptive drugs once   |  |  |
| 3.5 . 1                          |  |                    |   | the client knows she is pregnant.  |  |  |
| Mental<br>Health                 |  |                    |   | * A. No history of mental health diagnosis, but answers "Yes" to "In the last month, have you felt down, depressed or hopeless?" or showing potential symptoms of depression, but has negative score on standardized depression screening tool. i.e. Edinburgh, CES-D                    |  |  |
| Severe<br>Mental                 |  |                    |   | *B. History of mental health treatment but is stable, or history of postpartum depression with previous pregnancy, and negative score on standardized depression screening tool.   |  |  |
| Illness (SMI) and                |  |                    |   | *B Current mental health diagnosis and is engaged in mental health treatment   |  |  |
| Perinatal<br>Mood<br>Disorder    |  |                    |   | *C. Mental health symptoms of depression are evidenced by positive score on standardized depression screening tool   |  |  |
|                                  |  |                    |   | *C. Client has a mental health diagnosis and exhibiting active symptoms which are interfering with general functioning.  |  |  |
| Develop-<br>mental<br>Disability | *A. Severe developmental disability which could impact the woman's a herself during the pregnancy or an infant, but has adequate support |                    | *A. Severe developmental disability which could impact the woman's ability to take care of herself during the pregnancy or an infant, but has <b>adequate</b> support system, and <b>demonstrates</b> evidence of follow through with health care appointments/advice and self care |  |  |  |
| Disability                       |  |                    |   | *C. Severe developmental disability which impacts the woman's ability to take care of herself during the pregnancy or an infant and has an <b>inadequate</b> support system or <b>does not demonstrate</b> evidence of follow through with health care appointments/advice and self care |  |  |
| Intimate<br>Partner<br>Violence  |  |                    |   | <b>A.</b> In the last year, the woman's intimate partner or father of baby (FOB) has committed or threatened physical/sexual violence against her  |  |  |
| ☐ Check the box                  | to ackn  | owledg             | e all the N   | MSS targeted risk factors have been screened for and initial   |  |  |
| Screen Date                      |  | (                  | Complete  | ed by Level of service   |  |  |
| Screen Date                      | Completed by   |                    |   | ed by Level of service   |  |  |
| Screen Date                      |  | (                  | Complete  | ed by Level of service   |  |  |
|                                  | targetee<br>east one   | d risk fa<br>B and | actor or an<br>no C risk  | gnancy to two months post-pregnancy): ny As but no Bs or Cs criteria identified criteria   |  |  |

| Client Name: | DOB: | Client ID #: | 07/01/09 |
|--------------|------|--------------|----------|
|              |      |              |          |